

Dept. of Math

B.Sc. Part I Paper-I
Theory of Equations

Relation between roots and coefficients
of a polynomial equations

Let α, β, γ be the roots of $x^3 + px^2 + qx + r = 0$.
Then, writing the expression $x^3 + px^2 + qx + r$ in
the terms of α, β and γ gives $(x-\alpha)(x-\beta)(x-\gamma)$.

$$\begin{aligned} \therefore x^3 + px^2 + qx + r &= (x-\alpha)(x-\beta)(x-\gamma) \\ &= (x^2 - [\alpha+\beta]x + \alpha\beta)(x-\gamma) \\ &= x^3 - (\alpha+\beta)x^2 + \alpha\beta x - \gamma x^2 \\ &\quad + (\alpha+\beta)\gamma x - \alpha\beta\gamma \\ &= x^3 - (\alpha+\beta+\gamma)x^2 + (\alpha\beta+\beta\gamma+\gamma\alpha)x \\ &\quad - \alpha\beta\gamma \end{aligned}$$

From equating coefficients, we get

$$(a) \alpha + \beta + \gamma = -p \quad (b) \alpha\beta + \beta\gamma + \gamma\alpha = q \quad (c) \alpha\beta\gamma = -r.$$

This, of course, applies to a cubic equation.

Let us extend this to a more general equation

In general, if $\alpha_1, \alpha_2, \alpha_3, \dots, \alpha_n$ are the roots
of the equation

$$p_0 x^n + p_1 x^{n-1} + p_2 x^{n-2} + \dots + p_{n-1} x + p_n = 0$$

where $p_0 \neq 0$, then

(i) Sum of the roots = $-\frac{p_1}{p_0}$

(ii) Sum of products of the roots, three at a time = $-\frac{p_3}{p_0}$

(iii) Sum of products of the roots, n at a time
= $(-1)^n \frac{p_n}{p_0}$

The Factor theorem says that if a polynomial $P(x)$ has root r , then $x-r$ divides $P(x)$.

Since, polynomial with complex coefficients always have exactly the same number of roots as its degree (counting multiplicity), and they have unique factorization, that means that if $p(x)$ is the polynomial

$$x^n + p_1 x^{n-1} + \dots + p_{n-1} x + p_0$$

and r_1, r_2, \dots, r_n are roots of $p(x)$, then we can factor $p(x)$ as

$$P(x) = x^n + p_1 x^{n-1} + \dots + p_{n-1} x + p_0 \\ = (x-r_1)(x-r_2) \dots (x-r_n).$$

When multiply out $(x-r_1) \dots (x-r_n)$, and then put together the powers of x , we get a polynomial expression in which the coefficients are written in the terms of the roots. For example, to get the coefficient of x , note that we will get an x in the first binomial by each of the constants in the rest, giving us $(-1)^{n-1} (r_2 \dots r_n) x$.

Another when we multiply the x in the second binomial by each of the constants in the rest; another when you multiply the x in the third binomial by the constants in the rest; etc

if we work this out, we will find that when we expand $(x-r_1) \dots (x-r_n)$ and then group together the powers of x , you will have

- * The coefficient of x^n is 1
- * The coefficient of x^{n-1} is $-(r_1 + \dots + r_n)$.
- * The coefficient of x is $(-1)^{n-1} (r_1 r_2 \dots r_{n-1} + r_1 r_2 \dots r_{n-2} r_n + \dots + r_2 \dots r_n)$.
- * The constant coefficient is $(-1)^n (r_1 \dots r_n)$.

But for two polynomials to be equal they have to be equal coefficient by coefficient.

So that means that

$$P_1 = (-1)^1 (x_1 + \dots + x_n)$$

$$P_2 = (-1)^2 (x_1 x_2 + x_1 x_3 + \dots + x_1 x_n + x_2 x_3 + \dots + x_{n-1} x_n)$$

$$P_3 = (-1)^3 (x_1 x_2 x_3 + x_1 x_2 x_4 + \dots + x_1 x_2 x_n + x_1 x_3 x_4 + \dots + x_{n-2} x_{n-1} x_n)$$

⋮

$$P_n = (-1)^n (x_1 \dots x_n).$$

i.e.

- The sum of the roots is $-P_1$;
- The sum of all products of two roots is $(-1)^2 P_2$;
- The sum of all products of three roots is $(-1)^3 P_3$;
- ⋮
- The sum of all products of $n-1$ roots is $(-1)^{n-1} P_{n-1}$;
- The product of all roots is $(-1)^n P_n$.

How about a polynomial that is not monic,

$$P(x) = p_0 x^n + p_1 x^{n-1} + \dots + p_{n-1} x + p_n, \quad p_0 \neq 0?$$

Note that x is a root of $P(x)$ if and only if it is a root of

$$P(x) = \frac{1}{p_0} P(x) = x^n + \frac{p_1}{p_0} x^{n-1} + \dots + \frac{p_{n-1}}{p_0} x + \frac{p_n}{p_0}.$$

So the argument above applies of $P(x)$.