

number of molecules of

i.e

Derivation of gas laws

$PV = \text{Constant}$

or $P \propto \frac{1}{V}$ at constant Temperature T

this is Boyle's law

Derivation of Charles's Law from Kinetic Gas Equation

We know,

$PV = \frac{2}{3} RT$

or $V = \frac{2}{3} \cdot \frac{R \cdot T}{P}$

$V = \frac{2R}{3P} \cdot T$ ———

If Pressure P is constant,

then $V \propto T$

3. Avogadro's Law Derivation:

When two gases have the same Pressure and Volume,
then $P_1 V_1 = P_2 V_2$

We know from Kinetic gas eqⁿ $PV = \frac{1}{3} m n c^2$

Applying Kinetic gas eqⁿ

$\frac{1}{3} m_1 n_1 c_1^2 = \frac{1}{3} m_2 n_2 c_2^2$

$\frac{2}{3} \cdot \frac{1}{2} m_1 n_1 c_1^2 = \frac{2}{3} \cdot \frac{1}{2} m_2 n_2 c_2^2$

or $\frac{1}{2} m_1 n_1 c_1^2 = \frac{1}{2} m_2 n_2 c_2^2$ ——— (1)

When the temperature of these two gases is also the same,
then their Kinetic energy per mole will also be the same

i.e $\frac{1}{2} m_1 c_1^2 = \frac{1}{2} m_2 c_2^2$ ——— (2)

Dividing equation (1) by (2)

we get, $n_1 = n_2$

Thus the equal volumes of all gases under the same conditions of temperature and Pressure have the same number of moles.

This is Avogadro's Law.