

Pinky Rani
Assistant Professor (Guest Faculty)
Department of Economics
Maharaja College
Veer Kunwar Singh University, Ara
Class: B.A. Economics (Sem-2)
Paper: MJC-2
Topic: Rules of Differentiation

Product Rule of Differentiation

According to the product rule of derivatives, if the function $f(x)$ is the product of two functions $u(x)$ and $v(x)$, then the derivative of the function is given by:

If $f(x) = u(x) \times v(x)$, then:

$$f'(x) = u'(x) \times v(x) + u(x) \times v'(x)$$

Example: Find the derivative of $x^2(x+3)$.

Solution: As per the product rule of derivative, we know;

$$f'(x) = u'(x) \times v(x) + u(x) \times v'(x)$$

Here,

$$u(x) = x^2 \text{ and } v(x) = x+3$$

Therefore, on differentiating the given function, we get;

$$f'(x) = d/dx[x^2(x+3)]$$

$$f'(x) = d/dx(x^2)(x+3) + x^2 d/dx(x+3)$$

$$f'(x) = 2x(x+3) + x^2(1)$$

$$f'(x) = 2x^2 + 6x + x^2$$

$$f'(x) = 3x^2 + 6x$$

$$f'(x) = 3x(x+2)$$

Quotient Rule of Differentiation

If $f(x)$ is a function, which is equal to ratio of two functions $u(x)$ and $v(x)$ such that;

$$f(x) = u(x)/v(x)$$

Then, as per the quotient rule, the derivative of $f(x)$ is given by;

$$f'(x) = \frac{u'(x) \times v(x) - u(x) \times v'(x)}{(v(x))^2}$$

Example: Differentiate $f(x) = (x+2)^3/\sqrt{x}$

Solution: Given,

$$f(x) = (x+2)^3/\sqrt{x}$$

$$= (x+2)(x^2+4x+4)/\sqrt{x}$$

$$= [x^3+6x^2+12x+8]/x^{1/2}$$

$$= x^{-1/2}(x^3+6x^2+12x+8)$$

$$= x^{5/2}+6x^{3/2}+12x^{1/2}+8x^{-1/2}$$

Now, differentiating the given equation, we get;

$$f'(x) = 5/2x^{3/2} + 6(3/2x^{1/2})+12(1/2x^{-1/2})+8(-1/2x^{-3/2})$$

$$= 5/2x^{3/2} + 9x^{1/2} + 6x^{-1/2} - 4x^{-3/2}$$

Chain Rule of Differentiation

If a function $y = f(x) = g(u)$ and if $u = h(x)$, then the chain rule for differentiation is defined as;

$$dy/dx = (dy/du) \times (du/dx)$$

This rule is majorly used in the method of substitution where we can perform differentiation of composite functions.

Let's have a look at the examples given below for better understanding of the chain rule differentiation of functions.

Example 1:

Differentiate $f(x) = (x^4 - 1)^{50}$

Solution:

Given,

$$f(x) = (x^4 - 1)^{50}$$

Let $g(x) = x^4 - 1$ and $n = 50$

$$u(t) = t^{50}$$

Thus, $t = g(x) = x^4 - 1$

$$f(x) = u(g(x))$$

According to chain rule,

$$df/dx = (du/dt) \times (dt/dx)$$

Here,

$$du/dt = d/dt (t^{50}) = 50t^{49}$$

$$dt/dx = d/dx g(x)$$

$$= d/dx (x^4 - 1)$$

$$= 4x^3$$

$$\text{Thus, } df/dx = 50t^{49} \times (4x^3)$$

$$= 50(x^4 - 1)^{49} \times (4x^3)$$

$$= 200 x^3(x^4 - 1)^{49}$$