

Solution of Laplace's equation in spherical polar co-ordinates: spherical harmonics

The Laplace's equation

$$\nabla^2 U = 0$$

in spherical polar co-ordinates (r, θ, ϕ) takes the form

$$\frac{\partial}{\partial r} \left(r^2 \frac{\partial U}{\partial r} \right) + \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial U}{\partial \theta} \right) + \frac{1}{\sin^2 \theta} \frac{\partial^2 U}{\partial \phi^2} = 0 \quad \text{--- (1)}$$

From this equation it is obvious that U is the function of (r, θ, ϕ) ; therefore by the method of separation of variables its solution may be expressed as

$$U(r, \theta, \phi) = R \Theta \Phi = RS \quad \text{--- (2)}$$

where R is function of r only, Θ is the function of θ only and Φ is the function of ϕ only.

The function $S(\theta, \phi) = \Theta \Phi$ --- (3)

in the function of θ and ϕ and is called the surface harmonic. If ϕ is constant, then S is the function of θ only and is called the zonal surface harmonic.

2010	MAY 2010			AUGUST		
5	12	19	26	20	27	3
6	13	20	27	21	28	4
7	14	21	28	22	29	5
8	15	22	29	23	30	6
9	16	23	30	24	31	7
10	17	24	31	25		8
11	18	25		26		9
	19	26		27		10
	20	27		28		11
	21	28		29		12
	22	29		30		13
	23	30				14
	24					15
	25					16

Substituting $u = RS$ from (4) in (1) and

dividing by RS , we get

$$\frac{1}{R} \frac{\partial}{\partial r} \left(r^2 \frac{\partial R}{\partial r} \right) + \frac{1}{S \sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial S}{\partial \theta} \right) + \frac{1}{S \sin^2 \theta} \frac{\partial^2 S}{\partial \phi^2} =$$

$$\frac{1}{R} \frac{\partial}{\partial r} \left(r^2 \frac{\partial R}{\partial r} \right) = - \frac{1}{S \sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial S}{\partial \theta} \right) - \frac{1}{S \sin^2 \theta} \frac{\partial^2 S}{\partial \phi^2} \quad (4)$$

In this equation the left hand side is the function of r only and right hand side is the function of θ and ϕ , therefore each side must be equal to the same constant $n(n+1)$ (say), n being a constant,

$$\frac{1}{R} \frac{\partial}{\partial r} \left(r^2 \frac{\partial R}{\partial r} \right) = n(n+1)$$

$$r^2 \frac{\partial^2 R}{\partial r^2} + 2r \frac{\partial R}{\partial r} - n(n+1)R = 0 \quad (5)$$

$$- \frac{1}{S \sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial S}{\partial \theta} \right) - \frac{1}{S \sin^2 \theta} \frac{\partial^2 S}{\partial \phi^2} = n(n+1)$$

$$\frac{1}{S \sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial S}{\partial \theta} \right) + \frac{1}{S \sin^2 \theta} \frac{\partial^2 S}{\partial \phi^2} + n(n+1)S = 0$$

The solution of equation (5) is

$$R = A r^n + B r^{-n-1} \quad (7)$$

2016					MAY					2016					
Mon	31	3	10	17	24	Mon	7	14	21	28	Mon	7	14	21	28
Tue		4	11	18	25	Tue	8	15	22	29	Tue	8	15	22	29
Wed		5	12	19	26	Wed	9	16	23	30	Wed	9	16	23	30

If $S = S_n$ is the solution of equation (6) then the solutions of Laplace's equation in spherical polar co-ordinates is expressed as

$$U = RS = (A r^n + B r^{-n-1}) S_n \quad \text{--- (8)}$$

This solution is called the spherical harmonic. The subscript n on S_n signifies that the same value of n must be used in both terms of equation (8)

— + —