

Pinky Rani
 Assistant Professor (Guest Faculty)
 Department of Economics
 Maharaja College
 Veer Kunwar Singh University, Ara
 Class: B.A. Economics (Part-3)
 Paper: 07
 Topic: Index number

Construction of Price Index Numbers (Formula and Examples):
Construction of price index numbers through various methods can be understood with the help of the following examples:

1. Simple Aggregative Method:

In this method, the index number is equal to the sum of prices for the year for which index number is to be found divided by the sum of actual prices for the base year.

The formula for finding the index number through this method is as follows:

$$P_{01} = \frac{\Sigma P_1}{\Sigma P_0} \times 100$$

Where P_{01} Stands for the index number
 ΣP_1 Stands for the sum of the prices for the year for which index number is to be found :
 ΣP_0 Stands for the sum of prices for the base year.

Commodity	Prices in Base Year 1980 (in Rs.) P_0	Prices in current Year 1988 (in Rs.) P_1
A	10	20
B	15	25
C	40	60
D	25	40
Total	$\Sigma P_0 = 90$	$\Sigma P_1 = 145$

$$\text{Index Number } (P_{01}) = \frac{\Sigma P_1}{\Sigma P_0} \times 100 ; P_{01} = \frac{145}{90} \times 100 ; P_{01} = 161.11$$

2. Simple Average of Price Relatives Method:

In this method, the index number is equal to the sum of price relatives divided by the number of items and is calculated by using the following formula:

$$P_{01} = \frac{\Sigma R}{N}$$

Where ΣR stands for the sum of price relatives i. e. $R = \frac{P_1}{P_0} \times 100$ and

N stands for the number of items.

Example

Commodity P_0	Base Year Prices (in Rs.) P_1	Current year Prices (in Rs.)	Price Relatives $R = \frac{P_1}{P_0} \times 100$
A	10	20	$\frac{20}{10} \times 100 = 200.0$
B	15	25	$\frac{25}{15} \times 100 = 166.7$
C	40	60	$\frac{60}{40} \times 100 = 150.00$
D	25	40	$\frac{40}{25} \times 100 = 160.0$
$N = 4$			$\Sigma R = 676.7$

$$\text{Index Number } (p_{01}) = \frac{\Sigma R}{N}$$

$$P_{01} = \frac{676.7}{4} ; P_{01} = 169.2$$

3. Weighted Aggregative Method:

In this method, different weights are assigned to the items according to their relative importance. Weights used are the quantity weights. Many formulae have been developed to estimate index numbers on the basis of quantity weights.

Some of them are explained below:

(i) **Laspeyre's Formula.** In this formula, the quantities of base year are accepted as weights.

$$P_{01} = \frac{\sum P_1 q_0}{\sum P_0 q_0} \times 100$$

Where P_1 is the price in the current year ; P_0 is the price in the base year ; and q_0 is the quantity in the base year.

(ii) **Paasche's Formula.** In this formula, the quantities of the current year are accepted as weights.

$$P_{01} = \frac{\sum P_1 q_1}{\sum P_0 q_1} \times 100$$

Where q_1 is the quantity in the current year.

(iii) **Dorbish and Bowley's Formula.** Dorbish and Bowley's formula for estimating weighted index number is as follows :

$$P_{01} = \frac{\frac{\sum P_1 q_0}{\sum P_0 q_0} + \frac{\sum P_1 q_1}{\sum P_0 q_1}}{2} \times 100 \quad \text{or} \quad p_{01} = \frac{L + P}{2}$$

Where L is Laspeyre's index and P is paasche's Index.

(iv) **Fisher's Ideal Formula.** In this formula, the geometric mean of two indices (i.e., Laspeyre's Index and paasche's Index) is taken :

$$p_{01} = \sqrt{\frac{\sum P_1 q_0}{\sum P_0 q_0} \times \frac{\sum P_1 q_1}{\sum P_0 q_1}} \times 100 \quad \text{or} \quad P_{01} = \sqrt{L \times P} \times 100$$

where L is Lespeyre's Index and P is paasche's Index.

Example

Comm- odity	Base Year		Current Year		$P_0 q_0$	$P_1 q_0$	$P_0 q_1$	$P_1 q_1$
	P_0	q_0	P_1	q_1				
A	10	5	20	2	50	100	20	40
B	15	4	25	8	60	100	120	200
C	40	2	60	6	80	120	240	360
D	25	3	40	4	75	120	100	160
Total					265 $\sum P_0 q_0$	440 $\sum P_1 q_0$	480 $\sum P_0 q_1$	760 $\sum P_1 q_1$

(i) Laspeyre's Formula :

$$p_{01} = \frac{\sum P_1 q_0}{\sum P_0 q_0} \times 100$$

$$p_{01} = \frac{440}{265} \times 100 = 166.04$$

(ii) Paasche' Formula :

$$p_{01} = \frac{\sum P_1 q_1}{\sum P_0 q_1} \times 100$$

$$p_{01} = \frac{700}{480} \times 100 = 158.3$$

(iii) Dorbish and Bowley's Formula :

$$p_{01} = \frac{\frac{\sum P_1 q_0}{\sum P_0 q_0} + \frac{\sum P_1 q_1}{\sum P_0 q_1}}{2} \times 100 = 162.2$$

$$p_{01} = \frac{\frac{440}{265} + \frac{760}{480}}{2} \times 100 = 162$$

(iv) Fisher's Ideal Formula :

$$p_{01} = \sqrt{\frac{\sum P_1 q_0}{\sum P_0 q_0} \times \frac{\sum P_1 q_1}{\sum P_0 q_1}} \times 100$$

$$p_{01} = \sqrt{\frac{440}{265} \times \frac{760}{480}} \times 100 = 162.1$$

4. Weighted Average of Relatives Method:

In this method also different weights are used for the items according to their relative importance.

The price index number is found out with the help of the following formula:

$$P_{01} = \frac{\sum RW}{\sum W}$$

where $\sum W$ stands for the sum of weights of different commodities :

and $\sum R$ stands for the sum of price relatives.

Commodity	Weights W	Base Prices Year P_0	Current Year Prices P_1	Price Relatives $R = \frac{P_1}{P_0} \times 100$	RW
A	5	10	20	$20/10 \times 100 = 200.0$	1000.0
B	4	15	25	$25/15 \times 100 = 166.7$	666.8
C	2	40	60	$60/40 \times 100 = 150.0$	300.0
D	3	25	40	$40/25 \times 100 = 160.0$	480.0
Total	$\sum W=14$				$\sum RW = 2446.8$

$$\text{Index Number } (P_{01}) = \frac{\sum RW}{\sum W}$$

$$p_{01} = \frac{2446.8}{14} = 174.8$$
