

B.Sc. SEMESTER-I

MJC(T)/MIC(T)

Inorganic Chemistry

Atomic Structure and Chemical Bonding

Atomic Structure- Quantum Numbers and their Significance

Dr. Jasmine Singh

Assistant Professor

Department of Chemistry

Maharaja College(VKSU), Ara

Quantum numbers

(1) Each orbital in an atom is specified by a set of three quantum numbers (n, l, m) and each electron is designated by a set of four quantum numbers (n, l, m and s).

(2) Principle quantum number (n)

(i) It was proposed by *Bohr's* and denoted by ' n '.

(ii) It determines the average distance between electron and nucleus, means it is denoted the size of atom.

$$r = \frac{n^2}{Z} \times 0.529 \text{ \AA}$$

(iii) It determine the energy of the electron in an orbit where electron is present.

$$E = -\frac{Z^2}{n^2} \times 313.3 \text{ Kcal per mole}$$

(iv) The maximum number of an electron in an orbit represented by this quantum number as $2n^2$. No energy shell in atoms of known elements possess more than 32 electrons.

(v) It gives the information of orbit K, L, M, N -----.

(vi) The value of energy increases with the increasing value of n .

(vii) It represents the major energy shell or orbit to which the electron belongs.

(viii) Angular momentum can also be calculated using principle quantum number

$$mvr = \frac{nh}{2\pi}$$

(3) Azimuthal quantum number (l)

(i) Azimuthal quantum number is also known as angular quantum number. Proposed by *Sommerfeld* and denoted by ' l '.

(ii) It determines the number of sub shells or sublevels to which the electron belongs.

(iii) It tells about the shape of subshells.

(iv) It also expresses the energies of subshells $s < p < d < f$ (increasing energy).

(v) The value of $l = (n - 1)$ always where ' n ' is the number of principle shell.

(vi) Value of l	=	0	1	2	3.....(n-1)
Name of subshell	=	s	p	d	f
Shape of subshell	=	Spherical	Dumbbell	Double dumbbell	Complex

(vii) It represents the orbital angular momentum. Which is equal to? $\frac{h}{2\pi} \sqrt{l(l+1)}$

(viii) The maximum number of electrons in subshell $= 2(2l + 1)$

s - subshell $\rightarrow$ 2 electrons d - subshell $\rightarrow$ 10 electrons

p - subshell $\rightarrow$ 6 electrons f - subshell $\rightarrow$ 14 electrons.

(ix) For a given value of ' n ' the total value of ' l ' is always equal to the value of ' n '.


(x) The energy of any electron is depend on the value of n & l because total energy $= (n + l)$. The electron enters in that sub orbit whose $(n + l)$ value or the value of energy is less.

(4) Magnetic quantum number (m)

- (i) It was proposed by *Zeeman* and denoted by ' m '.
- (ii) It gives the number of permitted orientation of subshells.
- (iii) The value of m varies from $-l$ to $+l$ through zero.
- (iv) It tells about the splitting of spectral lines in the magnetic field i.e. this quantum number proved the Zeeman effect.
- (v) For a given value of ' n ' the total value of ' m ' is equal to n^2 .
- (vi) For a given value of ' l ' the total value of ' m ' is equal to $(2l + 1)$.
- (vii) *Degenerate orbitals* : Orbitals having the same energy are known as degenerate orbitals. e.g. for p subshell p_x p_y p_z
- (viii) The number of degenerate orbitals of s subshell $= 0$.

(5) Spin quantum numbers (s)

- (i) It was proposed by *Goldshmidt & Ulen Back* and denoted by the symbol of 's'.
- (ii) The value of 's' is $+1/2$ and $-1/2$, which signifies the spin or rotation or direction of electron on its axis during movement.
- (iii) The spin may be clockwise or anticlockwise.
- (iv) It represents the value of spin angular momentum is equal to $\frac{h}{2\pi} \sqrt{s(s+1)}$.
- (v) Maximum spin of an atom = $1/2 \times \text{number of unpaired electron}$.



- (vi) This quantum number is not the result of solution of Schrödinger equation as solved for *H*-atom.

Distribution of electrons among the quantum levels

n	l	m	s	Designation of orbitals	Electrons present	Total no. of electrons
1 (K shell)	0	0	$+1/2, -1/2$	1s	2	2
2 (L shell)	0	0	$+1/2, -1/2$	2s	2	8
		+1	$+1/2, -1/2$	2p	6	
		0	$+1/2, -1/2$			
		-1	$+1/2, -1/2$			
3 (M shell)	0	0	$+1/2, -1/2$	3s	2	18
		+1	$+1/2, -1/2$	3p	6	
		0	$+1/2, -1/2$			
		-1	$+1/2, -1/2$			
	2	+2	$+1/2, -1/2$	3d	10	
			$+1/2, -1/2$			
			$+1/2, -1/2$			
			$+1/2, -1/2$			
		0	$+1/2, -1/2$			
			$+1/2, -1/2$			
			$+1/2, -1/2$			
			$+1/2, -1/2$			

4(N shell)	0	0	$+1/2, -1/2$	4s	2	32
	1	+1	$+1/2, -1/2$	4p	6	
		0	$+1/2, -1/2$			
		-1	$+1/2, -1/2$			
	2	+2	$+1/2, -1/2$	4d	10	
		+1	$+1/2, -1/2$			
		0	$+1/2, -1/2$			
		-1	$+1/2, -1/2$			
		-2	$+1/2, -1/2$			
	3	+3		4f	14	
		+2				
		+1				
		+0				
		-1				
		-2	$+1/2, -1/2$			
		-3	$+1/2, -1/2$			
		$+1/2, -1/2$				
	$+1/2, -1/2$					
	$+1/2, -1/2$					
	$+1/2, -1/2$					
	$+1/2, -1/2$					
	$+1/2, -1/2$					
	$+1/2, -1/2$					